

# Transportability and Feasibility of Social Cognition and Interaction Training (SCIT) in Community Settings

David L. Roberts

*Yale University School of Medicine, New Haven, USA*

David L. Penn

*University of North Carolina, Chapel Hill, USA*

Daniella Labate

*The Bridge Inc., New York, USA*

Seth A. Margolis and Abram Sterne

*F.E.G.S. Health and Human Services System, New York, USA*

**Background:** Social Cognition and Interaction Training (SCIT) is a manual-based group intervention designed to improve social cognition in schizophrenia. Initial studies conducted by the developers of SCIT suggest that the intervention has promise in ameliorating social cognitive dysfunction in both inpatients and outpatients. **Aims:** The current study is a preliminary evaluation of SCIT in community samples. **Method:** An uncontrolled, pre-post design was used in this initial feasibility study. A collaborative research-clinical approach was employed to enable research evaluation while also meeting the administrative goals of participating clinics, and working within the constraints of real-world clinical practice. Transportability, acceptability, and feasibility of SCIT were evaluated in terms of pre- and post-treatment evaluations, client attendance data ( $N = 50$ ), and clinic administrators' decisions about whether to integrate SCIT into regular programming. Social-cognitive outcome measures assessed emotion perception, Theory of Mind, and attributional bias. **Results:** These support the transportability, acceptability, and feasibility of SCIT in community settings. SCIT has been integrated into routine practice at several test sites. Tentative support was found for improvement in emotion perception and Theory of Mind, but not attributional bias. **Conclusions:** SCIT may be a promising intervention for community agencies serving individuals with psychotic disorders who seek to improve their social functioning.

*Keywords:* Schizophrenia, social cognition, Theory of Mind, emotion perception, attributional bias, transportability.

---

Reprint requests to David L. Roberts, Yale University School of Medicine, Department of Psychiatry, CMHC, Room 38, 34 Park Street, New Haven, CT 06519, USA. E-mail: david.l.roberts@yale.edu

## Introduction

Social functioning deficits (e.g. social skill, community functioning) are a core feature of schizophrenia. They are a key diagnostic criterion for the disorder (DSM-IV; APA, 1994), precede illness onset (Davidson, Reichenberg, Rabinowitz, Weiser and Kaplan, 1999; Marenco and Weinberger, 2000), and are a strong predictor of outcome (Perlick, Stastny, Mattis and Teresi, 1992; Sullivan, Marder, Liberman, Donahoe and Mintz, 1990). These deficits are only minimally improved via medication (Bellack, Schooler and Marder, 2004), the frontline treatment for schizophrenia. Additionally, many of the prominent psychosocial interventions for schizophrenia (i.e. social skills training and neurocognitive remediation) exhibit limited generalizability beyond their targeted domains to clinically significant social outcomes (Kurtz and Mueser, 2008; Twamley, Jeste and Bellack, 2003). Cognitive-behavioral interventions traditionally have not targeted social functioning (Roberts, Penn, Cather, Otto and Goff, 2004); however there is emerging evidence that, if appropriately focused, they may confer social functioning benefits (Granholtm, Ben-Zeev and Link, in press; Wykes, Steel, Everitt and Tarrier, 2008).

Recently, social cognition has been identified as a potentially promising treatment target for improving social functioning in schizophrenia (Green, Olivier, Crawley, Penn and Silverstein, 2005; Penn, Corrigan, Bentall, Racenstein and Newman, 1997; Penn, Sanna and Roberts, 2008). Social cognition refers to a set of cognitive processes applied to the recognition, adaptive processing, and effective use of social cues in real-world situations (Brothers, 1990). Social cognition is a promising treatment target because it is related to social functioning outcomes (Couture, Penn and Roberts, 2006). Some evidence suggests that social cognition may be more strongly related to social functioning than is neurocognition (Pinkham and Penn, 2006; Pollice et al., 2002; Vauth, Rusch, Wirtz and Corrigan, 2004), while other research suggests that social cognition and neurocognition exert an interactive effect on social functioning (Bell, Tsang, Greig and Bryson, 2009; Brekke, Kay, Lee and Green, 2005).

Key domains of social cognition that have been found to be abnormal in schizophrenia include emotion perception, Theory of Mind (ToM), and attributional bias (Penn et al., 2008). Emotion perception is typically operationalized as one's ability to judge others' emotional states based on their facial expression or vocal tone. ToM refers to the ability to infer others' thoughts, beliefs or intentions indirectly, based on their utterances and behaviors, and in light of contextual clues; ToM is similar to "perspective taking". Attributional bias refers to patterned tendencies in explaining the causes of events in one's life. Individuals with persecutory delusions may exhibit a *hostile attributional bias* whereby they attribute negative events to the hostile intentions of others.

Social Cognition and Interaction Training (SCIT) is a manual-based, psychosocial treatment that is designed to improve social functioning in schizophrenia by way of enhanced social cognition. SCIT has shown feasibility and preliminary evidence of efficacy among both inpatient (Penn et al., 2005; Combs et al., 2007) and outpatient samples (Roberts and Penn, 2009).

The current paper reports on an open transportability trial of SCIT conducted within a network of community agencies unaffiliated with the developers of SCIT. In holding with stage models of treatment development (Onken, Blaine and Battjes, 1997), transportability trials of novel interventions are typically conducted after an intervention has demonstrated evidence

of efficacy in a controlled trial. However, bridging the gap between research findings and clinical practice has often proven difficult, and many interventions that demonstrate efficacy fail to receive appropriate transportability and effectiveness evaluation (Dobson and Hamilton, 2002; Westen, 2002). There are multiple reasons for this gap, including limited incentives for community stakeholders to participate in treatment research (Addis and Krasnow, 2000), the difficulty of modifying established treatment programming in community and hospital clinics, and the favoring of internal validity research over external validity research by funding agencies and academic publishing houses (Dobson and Hamilton, 2002). In sum, the traditional stage model of treatment development has encountered obstacles in producing treatment interventions that are internally as well as ecologically valid and transportable. In light of these obstacles, more recent models of treatment development have highlighted the benefits of integrating transportability and effectiveness studies earlier in the development process (e.g. Weisz, Chu and Polo, 2004).

The purpose of the current study was twofold: 1) To conduct a preliminary, open-trial evaluation of SCIT's transportability, feasibility, and acceptability in community settings prior to conducting a large-scale randomized, controlled trial; 2) To explore the utility of a model of research/clinical collaboration for facilitating the conduct of effectiveness research. Feasibility and acceptability assessment included feedback questionnaires completed by participating clients and group leaders, and attendance data. To assess the transportability of SCIT's active treatment components, and to provide treatment information useful to providers (Weisz et al., 2004), measures of emotion perception, Theory of Mind, and attributional bias were administered at pre- and post-treatment.

## Method

### *Treatment setting*

SCIT groups were conducted in New York City in collaboration with two healthcare service agencies, each of which operates a network of outpatient mental health services. Groups were conducted at three rehabilitation-oriented treatment centers for individuals with severe and persistent mental illnesses: (1) an intensive psychiatric rehabilitation program, which provides mental health, case-management, social, and vocational services in promotion of its clients' independent living; (2) a rehabilitation program located within a supported living facility for adults; (3) an outpatient mental health day program, which provides a spectrum of rehabilitation-oriented outpatient services with the mission of promoting independent living and self-sufficiency for its clients.

Across sites, SCIT was offered as an adjunct to routine care, and followed the manual-specified schedule of 20 to 24 sessions over approximately 5 months. Each group was led by two clinicians (three with Bachelor's degrees and three with Master's degrees). The clinicians had an average of 8.1 years of experience ( $SD = 7.3$ ) working with clients with severe mental illness.

### *Treatment intervention*

SCIT is a 20-session manual-based intervention. All sessions begin with structured check-ins designed to increase emotional self-awareness, and the relationship between participants'

emotions and social judgments. Next, homework is reviewed to bridge content from the previous week to the current session. In the first several sessions, this is followed by psychoeducation and discussion of social cognitive principles. The bulk of most core sessions is spent learning specific social cognitive strategies, and then utilizing these strategies to analyze social cognitive stimuli (photographs, specialized videos, or incidents from group members' lives). Strategy practice is structured in the form of games (including feedback about right and wrong answers) or as collaborative data-gathering, confidence-estimating, or problem-solving exercises. For example, participants learn that *jumping to conclusions* in social situations may lead to negative outcomes. They learn that one strategy to avoid jumping to conclusions is to separate social facts (e.g. she is looking this way) from guesses (e.g. she is thinking about me). Participants practice making this distinction in photographs and videos of social scenes. The final five sessions of SCIT are spent applying social cognitive strategies, such as separating facts from guesses, to interpersonal problems in participants' day-to-day lives. All sessions end with assignment of homework.

#### *Collaborative research approach*

The study was conducted by two of the SCIT treatment developers (DR and DP), located in North Carolina, along with administrators (AS) and clinicians (DL and SM) from three mental health treatment agencies in New York City. Based on documented obstacles to effectiveness research, the following collaboration principles were established: 1) Increase clinical and administrative stakeholders' incentives for participation; 2) Maximize research rigor *within* the resource limitations of participating clinical agencies; 3) Actively incorporate input from clinicians and administrators to enhance the feasibility and transportability of SCIT within community settings. These principles were addressed, respectively, as follows.

To increase clinicians' and administrators' incentives for participation, opportunities were provided for collaboration in the conduct, preparation, presentation, and write-up of research products related to the project. This was motivating for clinicians interested in developing as researchers. It was also motivating for administrators who had institutional commitments to increasing the empirical basis of their treatment offerings.

The primary resource limitations of participating agencies regarded group composition and outcome assessment. Regarding the former, it was deemed infeasible by clinical administrators to offer the SCIT intervention to only the subset of clients who met narrow research criteria (e.g. diagnostic homogeneity, exclusion based on current substance use). Therefore, inclusion criteria were relaxed. Regarding the latter, no research staff was available to provide participant assessments, nor did clinicians have time in their busy schedules to conduct individual research evaluations with participants. Therefore, standard social-cognitive outcome instruments were modified for use in group format, and were implemented by treating clinicians in the context of group sessions. Clinicians received coaching and supervision regarding test administration during weekly supervision calls.

Supervision calls were also used as a venue for clinicians to provide input regarding the feasibility of the intervention and the utility of the manual and bundled treatment materials. Clinicians' input was recorded and, after completion of the trial, was utilized to enhance the subsequent version of the treatment manual. It was expected that this approach would maximize the transportability of the version of the intervention that would be evaluated in the subsequent randomized controlled trial (currently underway).

### *Participants*

Participants were recruited by agency clinicians using the inclusion criteria from a previous efficacy trial (Roberts and Penn, 2009) as a guideline, in combination with the clinical goals of the program and the perceived treatment needs of individual clients. Specifically, clinicians prioritized recruitment of participants who met the following inclusion/exclusion criteria: (1) Diagnosis of schizophrenia or schizoaffective disorder; (2) Reading ability above third grade level; (3) Fluency in English; (4) No diagnosis of current substance abuse or dependence; (5) Difficulty with social cognition or paranoia, as indicated by clinician or staff consensus of difficulty with social interactions; (6) Aged between 18 and 65. All participants were receiving regular psychiatric treatment, consisting of case management, medication management, group treatment (i.e. psychoeducation and skill building) and, in some cases, vocational services.

Specific recruitment procedures differed across treatment sites, reflecting different institutional standards for group implementation. For example, one of the sites held an informational meeting at which the SCIT intervention was described. Clients had the opportunity to sign-up, and additional clients were approached based on clinicians' judgment. Thus, across sites, no standard data are available on the percentage of clients receiving information about SCIT who elected to participate. Individuals who elected to participate in SCIT were offered the opportunity to complete pre- and post-treatment research assessments as part of the current study. A subset of 50 group attendees elected to do so, and comprise the sample for the present study.

### *Measures*

*Demographic and clinical information.* Treating clinicians collected baseline data on age, gender, ethnicity, years of education, diagnosis, years with mental illness, and living status from participants' medical charts.

*Transportability, acceptability, and feasibility.* These domains were assessed using multiple methods. First, patients and clinicians were asked to complete evaluations of the SCIT intervention and materials. Using a 3-point scale (not helpful, helpful, very helpful), patients were asked to provide an overall rating of the group, as well as ratings of whether the group helped them in thinking about social situations and in interacting socially (see Table 2). Clinicians used a similar three-point scale to judge whether SCIT helped them to support their clients in improving their social cognition and in improving their social interactions (Table 3). Clinicians also rated the usefulness of the manual and materials on a similar three-point scale.

Second, pre- and post-treatment assessments of social cognition were conducted. As noted above, resources were not available to conduct extensive, one-on-one assessments with participants. Three widely-used social cognitive measures were modified for use in the current study. All three were produced in paper-and-pencil format, using enlarged font, and simplified printed instructions that were enhanced and clarified verbally as needed by group leaders. Measures were administered in group-format during the first three and final three meetings of the SCIT treatment groups. (Efforts were made to administer measures one-on-one for participants who missed any of the assessment meetings.) Because SCIT calls for groups to be co-facilitated by two clinicians, during assessment sessions one clinician presented assessment instructions and testing stimuli while the other clinician ensured that participants

understood the directions and assisted those with special needs (e.g. language, reading, or writing limitations).

*Emotion perception* was measured with the Face Emotion Identification Task (FEIT; Kerr and Neale, 1993). The FEIT is comprised of 19 photographs of faces expressing one of six basic emotions (happy, sad, angry, afraid, surprised, and ashamed). The participant's task is to determine which of the six emotions is being expressed by each face. Performance is indexed as the number of correct responses. The FEIT was modified by projecting the facial images onto a screen using an LCD projector.

*Theory of Mind* was measured with the Hinting task (Corcoran, Mercer and Frith, 1995). The Hinting task consists of 10 brief, written vignettes describing a social interaction between two characters that ends with one uttering a hint (e.g. "Gosh, these suitcases are heavy!"). The participant must infer what the character really meant by the hint (e.g. "Will you help me carry them?"). A correct inference receives 2 points. In the standard format, if the respondent is incorrect, a second, more obvious hint is provided (e.g. "I don't know if I can carry all three!") and, if correct at this point, the respondent receives 1 point. Incorrect answers receive 0 points. Scores on the Hinting task range from 0 to 20, with higher scores indicating better skills at inferring the desires of others. Because group administration made it impractical to provide the second hint to only a subset of respondents, all respondents were provided with the second hint and given the opportunity to change or add to their original answer based on this second hint.

*Attributional style* was measured with the Ambiguous Intentions Hostility Questionnaire-Ambiguous items (AIHQ-A; Combs et al., 2007). The AIHQ-A is comprised of five short, written, second-person vignettes describing negative interpersonal events with ambiguous causality. Each of the five vignettes is followed by a Hostility question (e.g. "Why did the other person do what s/he did?"), an Aggression question (e.g. "How would you respond?"), and a Blame question (e.g. "How much would you blame the person?"). Scores on each range from 0 to 5; higher scores indicate greater bias. Participants respond to Hostility and Aggression items with long-hand written answers. These responses were coded by a single, non-clinician rater, blind to pre/post status who had achieved adequate reliability ( $\alpha > .75$ ) with two other independent raters. Blame scores were derived from subject responses on Likert-type (0 to 5) scales.

SCIT's acceptability was further evaluated by examining group participants' attrition and attendance data. Finally, naturalistic follow-up with the providing clinics after termination of the joint research project provided information as to the administrators', clients', and clinicians' experience with the intervention.

*Fidelity and supervision.* Group facilitators read the SCIT treatment manual, attended a half-day workshop (conducted by DP and DR), and consulted with the treatment developers prior to initiating treatment. Facilitators participated in weekly supervision calls with DP and DR. No formal measure of treatment fidelity was administered.

#### *Data analysis plan*

Due to the small sample size and preliminary nature of this study, visual inspection of cell counts was used to assess responses on the post-treatment evaluation measures.

**Table 1.** Demographic information ( $N = 50$ )

	Mean /%	<i>SD</i>
Age (years)	53.1	11.8
Female (%)	55.0	
Ethnicity (%)		
African Am.	42.9	
Caucasian	30.6	
Hispanic	22.4	
Years of education	12.4	3.4
Years with illness*	22.2	14.5
Chart diagnosis (%)		
Schizophrenia	40.8	
Schizoaffective	20.4	
Bipolar d/o	14.3	
Depressive disorder	12.2	
Other	12.2	
Living status (%)		
Independent	10.2	
MH supported	34.7	
Group home	55.1	

MH supported = Apartment with functional supports from a mental health provider.

\* Four participants identified as “10 +” are not included.

Social cognitive outcome data were evaluated by conducting a repeated measures MANOVA on the primary social cognitive variables (FEIT, Hinting task, and AIHQ hostility bias). Follow-up paired-samples *t*-tests were used to explore the statistical significance of pre-to-posttest change on these three variables individually, as well as on the other two AIHQ variables (aggression bias and blame score). Within-group effect sizes were calculated to estimate the magnitude of change from pre- to posttest. Cohen's *d* (Cohen, 1988) was calculated using Dunlap and colleagues' conservative calculation, which corrects for effect size inflation due to within-variable correlation in paired samples (Dunlap, Cortina, Vaslow and Burke, 1996).

## Results

Demographic and clinical characteristics of the sample are summarized in Table 1.

### *Attrition and attendance*

Of the 50 participants who completed baseline assessments, 38 completed the SCIT training, yielding a 24% drop-out rate. Of the 12 drop-outs, 4 occurred after initial assessment but prior to SCIT treatment, 7 occurred during the first three sessions of treatment (one due to physical illness), and one occurred later in treatment.

**Table 2.** Patient feedback regarding SCIT ( $N = 24$ )

Item	Not helpful $N$	Helpful $N$	Very helpful $N$
How useful was the group to you?	2	10	12
How much did the group help you in thinking about social situations?	2	15	7
How much did the group help you in the way you relate to other people?	2	12	10

**Table 3.** Group leader feedback regarding SCIT ( $N = 7$ )

Item	Not helpful $N$	Helpful $N$	Very helpful $N$
How helpful was the SCIT manual?	0	0	7
How much did SCIT help you support your clients in improving their social cognition?	0	4	3
How much did SCIT help you support your clients in improving their social interactions?	0	2	5

Attendance data were collected from 31 of the 38 study completers.<sup>1</sup> These participants attended a mean of 15.5 sessions ( $SD = 2.8$ ) out of a total of 20 to 23 sessions, yielding an average attendance rate of 69%. (The total number of sessions used to complete the manual differed slightly across cohorts.)

#### *Post-treatment evaluations*

Twenty-four participants completed post-treatment evaluations.<sup>2</sup> As shown in Table 2, the large majority rated SCIT as either “helpful” or “very helpful” in all three domains. Seven group leaders completed post-treatment evaluations. As shown in Table 3, the majority gave positive ratings for the usefulness of the SCIT manual and materials, and in supporting improvement of their clients’ social cognition and social interactions. No clinicians rated SCIT as “not helpful” on any of the three items.

#### *Integration of SCIT into regular programming*

Follow-up with agency administrators after the termination of the research project revealed that two of the three participating clinics continued providing SCIT independent of collaboration with the treatment developers. The third clinic reported that they did not continue the group because of staffing shortages. However, the parent agency arranged for training of staff at affiliated clinics, and has now incorporated SCIT into routine programming at two additional sites.

<sup>1</sup>Attendance data were not collected by leaders of one treatment group.

<sup>2</sup>The evaluation was completed by fewer participants than post-test assessments because it was offered optionally following completion of the treatment.

**Table 4.** Social cognitive treatment outcomes

Variable (N)	Pretest <i>M</i> ( <i>SD</i> )	Posttest <i>M</i> ( <i>SD</i> )	<i>d</i>
FEIT (32)*	9.81 (3.25)	10.84 (2.95)	0.33
Hinting task (34)**	13.18 (4.20)	14.97 (4.16)	0.43
AIHQ Hostility (32)	1.95 (0.62)	1.86 (0.60)	0.15
AIHQ Aggression (32)	1.90 (0.39)	1.88 (0.34)	0.06
AIHQ Blame (31)	2.81 (0.88)	2.68 (0.76)	0.16

\* $p < .05$ ; \*\* $p < .005$ .

*M* = Mean; *SD* = Standard Deviation; *d* = Effect size.

*Note:* *N* differences across measures are due to several participants' failure to attend a portion of the testing sessions or failure to complete measures in session.

### *Treatment outcome findings*

Of the 38 participants who completed the SCIT training, 34 completed post-treatment assessments. Reasons for not completing post-treatment assessment were absence on the days of assessment (3) and refusal (1). Within the full sample of 50, comparison of baseline social-cognitive performance among the 34 post-test completers and the 16 non-completers revealed no significant differences on any social-cognitive variables.

Table 4 summarizes pre- and post-test social cognitive performance. The omnibus repeated measures MANOVA conducted on the primary social cognitive variables was statistically significant ( $F(1,30) = 4.42$ ;  $p = .011$ ), indicating an overall change in social cognition from pre- to post-test. In follow-up paired-samples *t*-tests, participants showed a statistically significant improvement in FEIT (emotion perception) performance ( $t = 2.22$ ;  $p = .034$ ), and Hinting task (ToM) performance ( $t = 3.24$ ;  $p = .003$ ), which correspond to effect sizes in the small-medium range. No change was observed in AIHQ hostility bias, nor AIHQ Aggression bias or Blame score.

## **Discussion**

This study evaluated the transportability, feasibility, and acceptability of SCIT, a social cognitive intervention for schizophrenia, in community mental health settings. Results indicate that participating clinicians and clients generally found SCIT to be acceptable and perceived it favorably. Treatment-outcome findings suggest that SCIT may have conferred social-cognitive benefits to participating clients. Findings are discussed in detail below.

Data concerning the transportability and feasibility of SCIT were collected from multiple sources, and were generally promising. Post-treatment evaluation data from participating clients was quite positive, indicating that they found it to be acceptable and useful. Positive client feedback is consistent with similar evaluations completed by participants in previous SCIT treatment groups (Penn, Roberts, Combs and Sterne, 2007). This finding is also consistent with an emphasis on maximizing client engagement, which guided development of SCIT. Additionally, the attendance rate of approximately 69% is on par with the attendance rate achieved in our previous outpatient trial of SCIT (Roberts and Penn, 2009). The drop-out rate of 24% was higher than observed in this previous outpatient trial; however, it is on par with rates observed in previous outpatient trials of CBT for psychosis (Wykes et al., 2008), as

well as rates of disengagement from community treatment programs (Kreyenbuhl, Nossel and Dixon, 2009).

Clinician evaluations also supported the feasibility of SCIT. Specifically, all seven clinicians gave the treatment materials the most favorable rating of “very helpful”. This finding is promising in light of previous evidence that practicing clinicians often perceive standardized treatment manuals negatively (Addis and Krasnow, 2000). As noted above, clinicians’ input regarding the treatment was also solicited and recorded during weekly supervision calls throughout the study period. Most input consisted of two types: (1) Ideas for supplementary materials, such as additional homework options and increased incorporation of photographs and imagery; (2) Requests for suggestions on problem solving around unusual client presentations or difficult situations. (These suggestions were incorporated into the manual.) In light of previous research on the transportability of manualized interventions (Addis and Krasnow, 2000), it is notable that clinicians did not indicate that implementing SCIT placed undo burden on them in the context of their busy clinical schedules, did not request streamlining of the manual, and did not indicate that the standardized treatment was inappropriate for their heterogeneous client population.

Perhaps the most meaningful feasibility data is the fact that clinic administrators have incorporated SCIT into routine programming at an expanded set of clinics. This promising finding may be due in part to the fact that SCIT was initially developed within a clinical treatment context (Penn et al., 2007). Thus, consistent with the Deployment-Focused Model of treatment development (Weisz et al., 2004), SCIT was developed to employ generic elements that are common to typical inpatient and outpatient treatment settings, including group structure, weekly one-hour meetings, 5-month duration, psychoeducational and exercise based intervention, low demand for costly materials and technology, and minimal requirement for specialized staff training beyond general principles of cognitive, behavioral, and group intervention approaches.

Data collected regarding SCIT’s effectiveness must be interpreted minimally and with caution because of the methodological limitations of this study. SCIT was associated with statistically significant improvement in emotion perception. The effect size was attenuated relative to previous studies of SCIT conducted by the developers of the treatment (Combs et al., 2007; Roberts and Penn, 2009). Such attenuation is not uncommon when comparing the effectiveness of an intervention to its efficacy (e.g. Curtis, Ronin and Borduin, 2004), and may be influenced by a selection artifact driven by the increased heterogeneity and complexity of illness among community patients versus carefully-screened research participants (Westen and Morrison, 2001).

Participants improved significantly on Theory of Mind performance, replicating findings from previous inpatient studies of SCIT (Combs et al., 2007; Penn et al., 2005), but contrasting with the two previous outpatient trials that used SCIT intervention techniques (Horan et al., 2009; Roberts and Penn, 2009). In the context of previous results, it is possible that the low functional level of the current participants prevented a ceiling effect on the Hinting task, enabling improvement that was not possible in previous, higher functioning samples.

As in both previous outpatient studies that used SCIT intervention techniques (Horan et al., 2009; Roberts and Penn, 2009), participants’ pre- and posttest scores on all three attributional bias scales (hostility, aggression, and blame) were in the low-normal range (cf. Combs et al., 2007), rendering moot the possibility of meaningful decrease in this domain. As noted previously (Roberts and Penn, 2009), this suggests a floor effect such that SCIT

participants actually endorsed *less biased* judgments than non-ill controls. Given the risk of self-presentation effects on measures of hostility bias, future research in this area may do well to follow the lead of social psychology in using implicit or non-obvious measures (cf. Greenwald and Banaji, 1995). We recently developed the Mental State Inference Questionnaire with this aim in mind (Roberts, Fiszdon, DeGeorge and Tek, 2009).

No standard measures of social functioning improvement were administered, although this domain was assessed informally in both client and clinician feedback questionnaires (Tables 2 and 3). Clinicians gave SCIT particularly strong endorsement in this domain, while client endorsement was also positive. Because social functioning improvement is the most important SCIT outcome domain, it will be crucial to formally assess this domain in future effectiveness research on SCIT.

This study has several notable limitations. It is a small, uncontrolled trial that used a convenience sample. Additionally, assessments were administered (1) by the treating clinicians, (2) in group format, and (3) using modified materials. All three of these factors threaten the internal validity of assessment results, and the first two could have inflated Type I error (Tarrier and Wykes, 2004). Additionally, the data used in this study are from the self-selected subset of patients who participated in both the pre- and post-test evaluations. It is possible that this subset found the curriculum more relevant and appealing than the subset of individuals who dropped out of the group or declined to participate in assessments. However, this possibility may be mitigated by the fact that all but one drop-out occurred prior to the fourth SCIT session. Thus, it could be argued that participants who gave SCIT a chance generally completed the intervention. Nonetheless, client feedback data may be an overestimation of the subjectively perceived helpfulness of the group to the broader population of outpatients, and it remains possible that this self-selection inflated treatment outcome findings.

These threats were known at the outset, and are largely an artifact of limited agency resources. The general convergence of the current outcome findings with previous research on SCIT strengthens our confidence in the validity of the data. Nonetheless, the current findings must be interpreted cautiously. As the emphasis on effectiveness research in mental health grows, there will be continued need for novel methods of maximizing assessment validity while minimizing agency cost and client burden. In this light, perhaps the most promising finding from the current study is the success of the research-clinical collaboration approach that was used and the dissemination of a promising treatment intervention that this collaboration generated.

### Acknowledgements

We wish to thank Laurel Carpenter, Michael Cavallaro, Amy Dorin, Ellen Stoller and Marjorie Stuckle, as well as the employees of F.E.G.S. Health and Human Services System and The Bridge, Inc. for their contributions to this project.

### References

- Addis, M. E. and Krasnow, A. D.** (2000). A national survey of practicing psychologists' attitudes toward psychotherapy treatment manuals. *Journal of Consulting and Clinical Psychology*, 68, 331–339.
- American Psychiatric Association** (1994). *Diagnostic and Statistical Manual of Mental Disorders* (4th ed.). Washington, DC: Author.

- Brothers, L.** (1990). The social brain: a project for integrating primate behavior and neurophysiology in a new domain. *Concepts in Neuroscience, 1*, 27–61.
- Bell, M., Tsang, H. W., Greig, T. C. and Bryson, G. J.** (2009). Neurocognition, social cognition, perceived social discomfort, and vocational outcomes in schizophrenia. *Schizophrenia Bulletin, 35*, 738–747.
- Bellack, A. S., Schooler, N. and Marder, S. R.** (2004). Do Clozapine and Risperidone affect social competence and problem solving? *American Journal of Psychiatry, 161*, 364–367.
- Brekke, J., Kay, D. D., Lee, K. S. and Green, M. F.** (2005). Biosocial pathways to functional outcome in schizophrenia. *Schizophrenia Research, 80*, 213–225.
- Cohen, J.** (1988). *Statistical Power Analysis for the Behavioral Sciences*. Hillsdale, NJ: Erlbaum.
- Combs, D. R., Adams, S. D., Penn, D. L., Roberts, D. L., Tiegreen, J. and Stem, P.** (2007). Social Cognition and Interaction Training for inpatients with schizophrenia spectrum disorders: preliminary findings. *Schizophrenia Research, 91*, 112–116.
- Combs, D. R., Penn, D. L., Wicher, M. and Waldheter, E.** (2007). The Ambiguous Intentions Hostility Questionnaire (AIHQ): a new measure for evaluating attributional biases in paranoia. *Cognitive Neuropsychiatry, 12*, 128–143.
- Corcoran, R., Mercer, G. and Frith, C.** (1995). Schizophrenia, symptomatology and social inference: investigating “theory of mind” in people with schizophrenia. *Schizophrenia Research, 17*, 5–13.
- Couture, S. M., Penn, D. L. and Roberts, D. L.** (2006). The functional significance of social cognition in schizophrenia: a review. *Schizophrenia Bulletin, 32*, S44–S63.
- Curtis, N. M., Ronan, K. R. and Borduin, C. M.** (2004) Multisystemic treatment: a meta-analysis of outcome studies. *Journal of Family Psychology, 18*, 411–419.
- Davidson, M., Reichenberg, A., Rabinowitz, J., Weiser, M. and Kaplan, Z.** (1999). Behavioral and intellectual markers for schizophrenia in apparently healthy male adolescents. *American Journal of Psychiatry, 156*, 1328–1335.
- Dobson, K. S. and Hamilton, K. E.** (2002). The stage model for psychotherapy manual development: a valuable tool for promoting evidence-based practice. *Clinical Psychology: Science and Practice, 9*, 407–409.
- Dunlap, W. P., Cortina, J. M., Vaslow, J. B. and Burke, M. J.** (1996). Meta-analysis of experiments with matched groups or repeated measures designs. *Psychological Methods, 1*, 170–177.
- Granholm, E., Ben-Zeev, D. and Link, P. C.** (in press). Social disinterest attitudes and group cognitive-behavioral social skills training for functional disability in schizophrenia. *Schizophrenia Bulletin*.
- Green, M. F., Olivier, B., Crawley, J. N., Penn, D. L. and Silverstein, S.** (2005). Social cognition in schizophrenia: recommendations from the MATRICS new approaches conference. *Schizophrenia Bulletin, 31*, 882–887.
- Greenwald, A. G. and Banaji, M. R.** (1995). Implicit social cognition: attitudes, self-esteem, and stereotypes. *Psychological Review, 102*, 4–27.
- Horan, W. P., Kern, R. S., Shokat-Fadai, K., Sergi, M. J., Wynn, J. K. and Green, M. F.** (2009). Social cognitive skills training in schizophrenia: an initial efficacy study of stabilized outpatients. *Schizophrenia Research, 107*, 47–54.
- Kerr, S. L. and Neale, J. M.** (1993). Emotion perception in schizophrenia: specific deficit or further evidence of generalized poor performance? *Journal of Abnormal Psychology, 102*, 312–318.
- Kreyenbuhl, J., Nossel, H. R. and Dixon, L. B.** (2009). Disengagement from mental health treatment among individuals with schizophrenia and strategies for facilitating connections to care: a review of the literature. *Schizophrenia Bulletin, 35*, 696–703.
- Kurtz, M. M. and Mueser, K. T.** (2008). A meta-analysis of controlled research on social skills training for schizophrenia. *Journal of Consulting and Clinical Psychology, 76*, 491–504.
- Marengo, S. and Weinberger, D. R.** (2000). The neurodevelopmental hypothesis of schizophrenia: following a trail of evidence from cradle to grave. *Development and Psychopathology, 12*, 501–527.

- Onken, L. S., Blaine, J. D. and Battjes, R. J.** (1997). Behavioral therapy research: a conceptualization of a process. In S. W. Henggeler and A. B. Santos (Eds.), *Innovative Approaches for Difficult-to-Treat Populations* (pp. 477–485). Washington, DC: American Psychiatric Press.
- Penn, D. L., Corrigan, P. W., Bentall, R., Racenstein, J. M. and Newman, L.** (1997). Social cognition in schizophrenia. *Psychological Bulletin*, *121*, 114–132.
- Penn, D. L., Roberts, D. L., Combs, D. and Sterne, A.** (2007). The development of the Social Cognition and Interaction Training program for schizophrenia spectrum disorders. *Psychiatric Services*, *58*, 449–451.
- Penn, D. L., Roberts, D., Munt, E. D., Silverstein, E., Jones, N. and Sheitman, B.** (2005). A pilot study of social cognition and interaction training (SCIT) for schizophrenia. *Schizophrenia Research*, *80*, 357–359.
- Penn, D. L., Sanna, L. and Roberts, D. L.** (2008). Social cognition in schizophrenia: an overview. *Schizophrenia Bulletin*, *34*, 408–411.
- Perlick, D., Stastny, P., Mattis, S. and Teresi, J.** (1992). Contribution of family, cognitive, and clinical dimensions to long-term outcome in schizophrenia. *Schizophrenia Research*, *6*, 257–265.
- Pollice, R., Roncone, R., Falloon, I. R. H., Mazza, M., DeRisio, A., Necozone, S., Morosini, P. and Casacchia, M.** (2002). Is theory of mind in schizophrenia more strongly associated with clinical and social functioning than with neurocognitive deficits? *Psychopathology*, *35*, 280–288.
- Pinkham, A. E. and Penn, D. L.** (2006). Neurocognitive and social cognitive predictors of interpersonal skill in schizophrenia. *Psychiatry Research*, *143*, 167–178.
- Roberts, D. L., Fiszdton, J. M., DeGeorge, P. and Tek, C.** (2009). Impression-management effects in paranoia assessment. *Schizophrenia Bulletin*, *35*, Suppl, 2–3.
- Roberts, D. L. and Penn, D. L.** (2009). Social Cognition and Interaction Training (SCIT) for outpatients with schizophrenia: a preliminary study. *Psychiatry Research*, *166*, 141–147.
- Roberts, D. L., Penn, D. L., Cather, C., Otto, M. and Goff, D. C.** (2004). Should CBT target the social impairments associated with schizophrenia? *Journal of Cognitive Psychotherapy: An International Quarterly*, *18*, 255–264.
- Sullivan, G., Marder, S. R., Liberman, R. P., Donahoe, C. P. and Mintz, J.** (1990). Social skills and relapse history in outpatient schizophrenics. *Psychiatry*, *53*, 340–345.
- Tarrier, N. and Wykes, T.** (2004). Is there evidence that cognitive behaviour therapy is an effective treatment for schizophrenia? A cautious or cautionary tale? *Behaviour Research and Therapy*, *42*, 1377–1401.
- Twamley, E. W., Jeste, D. V. and Bellack, A. S.** (2003). A review of cognitive training in schizophrenia. *Schizophrenia Bulletin*, *29*, 359–382.
- Vauth, R., Rusch, N., Wirtz, M. and Corrigan, P. W.** (2004). Does social cognition influence the relation between neurocognitive deficits and vocational functioning in schizophrenia? *Psychiatry Research*, *128*, 155–165.
- Weisz, J. R., Chu, B. C. and Polo, A. J.** (2004). Treatment dissemination and evidence-based practice: strengthening intervention through clinician-researcher collaboration. *Clinical Psychology: Science and Practice*, *11*, 300–307.
- Westen, D.** (2002). Manualizing manual development. *Clinical Psychology: Science and Practice*, *9*, 416–418.
- Westen, D. and Morrison, K.** (2001). A multidimensional meta-analysis of treatments for depression, panic, and generalized anxiety disorder: an empirical examination of the status of empirically supported therapies. *Journal of Consulting and Clinical Psychology*, *6*, 875–899.
- Wykes, T., Steel, C., Everitt, B. and Tarrier, N.** (2008). Cognitive behavior therapy for schizophrenia: effect sizes, clinical models, and methodological rigor. *Schizophrenia Bulletin*, *34*, 523–537.